AD-A117 051

BROWN UNIV PROVIDENCE RI LEFSCHETZ CENTER FOR DYNAMI--ETC F/6 12/1

STABLE EQUILIBRIA WITH VARIABLE DIFFUSION. (U)

MAR 82 M CHIPOT, J K HALE


LCOS-M-82-5

AFOSR-TR-82-0511

END

OR-322

OR-322





STABLE EQUILIBRIA WITH VARIABLE DIFFUSION

by

Michel Chipot

Visitor to:
Lefschetz Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, Rhode Island 02912

STIC ELECTE JUL 1 9 1982

F

On leave from:
Universite de Nancy I - UER.
Sc. Mathematiques
Case off. 239-54506 Vandoeuvre les Nancy-Cedex, France

and .

Jack K. Hale

Lefschetz Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, Rhode Island 02912

March 21, 1982

LCDS - #M-82-5

- This research has been supported in part by the Air Force Office of Scientific Research under contract ## -AFOSR 81-0198.
 - This research has been supported in part by the Air Force Office of Scientific Research under contract #AF-AFOSR 81-0198, in part by the National Science Foundation under contract #MCS 79-05774-05 and in part by the U. S. Army Research Office under contract #ARO-DAAG-29-79-C-0161.

Approved for public release; distribution unlimited.

82 07 19 135

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM	
AFOR-TR- 8 2 0 5 11 2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitio)	5. TYPE OF REPORT & PERIOD COVERED	
STABLE EQUILIBRIA WITH VARIABLE DIFFUSION	INTERIM	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s)	8 CONTRACT OR GRANT NUMBER(a)	
Michel Chipot Jack K. Hale	AFOSR-81-0138	
PERFORMING ORGANIZATION NAME AND ADDRESS Lefschetz Center for Dynamical Systems	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Division of Applied Mathematics Brown University, Providence, Rhode Island 02912	61102F 2304/A4	
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
AFOSR/NM Bolling AFB, DC 20332	MAR 1982 13. NUMBER OF PAGES	
	7	
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)	15. SECURITY CLASS. (of this report)	
	UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED		
17. DISTRIBUTION STATEMENT (of the ebetract entered in Block 20, if different from	m Report)	
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
For a scalar nonlinear parabolic equation in one space dimension with homogeneous Neumann boundary conditions, criteria are given on the diffusion coefficient to ensure that the stable equilibrium solutions are constant functions regardless of the nonlinearities. The Dirichlet problem is also discussed.		

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)		
		1
·		- 1
	•	J
		- 1
	•	- 1
	<u>-</u>	1
		1
	• •	1
•		Ì
		ļ
		I
		l l
		1
		i
		ı
		ļ
		1
		- }
		i
		l l
		1
		1
		l
		- 1
		j
		- 1
		- 1
		1
		1
		}
		- [
		1
		•
•		
		1
•		- 1
	•	- 1
	•	
		Í

Stable equilibria with variable diffusion

bу

Michel Chipot and Jack K. Hale

ABSTRACT

For a scalar nonlinear parabolic equation in one space dimension with homogeneous Neumann boundary conditions, criteria are given on the diffusion coefficient to ensure that the stable equilibrium solutions are constant functions regardless of the nonlinearities. The Dirichlet problem is also discussed.

	ı
Accession For	l
. 3 GRANI	l
TAB 🔲	
iannounced [1
	1
Jantification]
and the second s	}
BV	4
Distribution/	1
	7
ivnilability Codes	4
Avail and/or	ı
Trist Special	į
	1
11	1
111	1
V ' \	_
The second secon	_

DTIQ

COPY

AIR FORCE OFFICE OF SCIENTIFIC RESEARCY (AFSC)
NOTICE OF TO MSWITTAL TO DTIC
This technical inductions have reviewed end is
approved to the science law AFR 190-12.
Distribute to the load.
MATTHEW J. KELLIN
Chief, Technical Information Division

Consider the equation

(1)
$$u_t = (au_x)_x + f(u), \quad 0 < x < 1,$$

with homogeneous Neumann conditions

(2)
$$u_x = 0$$
 at $x = 0$, $x = 1$,

where $a \in C^2[0,1]$ is a positive function and $f \in C^2(\mathbb{R})$. System (1), (2) defines a local dynamical system in $H^1(0,1) = W^{1,2}(0,1)$ (see Henry [8]). Furthermore, the ω -limit set of any bounded orbit is exactly one equilibrium point; that is, a solution of the equations

(3)
$$(au_x)_x + f(u) = 0, \quad 0 < x < 1$$

(4)
$$u_x = 0$$
 at $x = 0$, $x = 1$

(see Matano [11], Hale and Massatt [6], Zelenyak [15]).

The first result is the following.

Theorem 1. If $a''(x) \le 0$ on [0,1], then every nonconstant equilibrium solution of (1),(2) is unstable.

For a(x) = c > 0, where c is a constant, this result was proved by Chafee [2]. Note that the conclusion in the theorem is valid for every function f. Matano [13] has given an example (1),(2) for which there are stable non-constant equilibrium solutions. The function f can be chosen to be a cubic with three simple zeros and the function a is ≥ 1 on intervals $[0,\alpha]$, $[\beta,1]$ and $\leq \epsilon$ on $[\gamma,\delta]$, $\alpha < \gamma < \delta < \beta$, and ϵ is sufficiently small. Numerical examples indicated the same property can occur with $a \geq 1$ on $[0,\alpha]$ and $\leq \epsilon$ on

 $[\beta,1]$ with $\alpha < \beta$. These examples show clearly that more work is needed to determine the optimal conditions on a for the conclusions of the theorem to hold.

For a more general parabolic equation in several space dimensions in a bounded domain Ω and the diffusion coefficients constant, Casten and Holland [1], Matano [12] have shown that Theorem 1 is true if Ω is convex. The authors have had no success in extending this result in a significant way with variable diffusion coefficients. If Ω is not convex and the diffusion coefficients are constants, nonconstant equilibrium solutions can occur (see Matano [12], Hale and Vegas [7], Vegas [14]) for certain nonlinear functions f. If the diffusion coefficient in (1) is constant and f is a function of $u(\cdot,t)$, for example, $f(u(x,t)) = g(u(x,t), \int_0^1 \alpha(x)u(x,t)dx)$, then stable nonequilibrium solutions can occur (see Chafee [3]). A similar situation was considered in several space dimensions and nonconvex domains by Keyfitz and Kuiper [9]. For constant diffusion in (1) and f replaced by $\dot{s}(x)f(u)$, Fleming [5], (see also Henry [8]) has considered how the existence of stable nonconstant solutions depend on s(x). All of these papers should be reconsidered with variable diffusion.

There is an analogue of Theorem 1 for the Dirichlet problem.

Theorem 2. Consider the equation (1) with Dirichlet boundary conditions

$$u = 0$$
 at $x = 0$, $x = 1$.

If $a''(x) \le 0$ on [0,1] and v(x) is a nonconstant equilibrium solution such that $v_x = 0$ at two points in (0,1), then $v_x = 0$ is unstable.

For a(x) = c > 0, this result was proved by Chafee and Infante [4], Maginu [10].

Proof of Theorem 1. If v is a solution of (3), (4) and

$$\mathcal{L}(\phi) = \int_0^1 [a\phi_x^2 - f'(v)\phi^2] dx,$$

then the first eigenvalue λ_1 of the operator

$$Lu = -(au_x)_x - f'(v)u$$

is given by

$$\lambda_1 = \min\{\mathcal{U}(\phi) : \phi \in H^1(0,1), |\phi|_{L^2(0,1)} = 1\}.$$

Furthermore, the equilibrium solution v of (1),(2) is unstable if $\lambda_1 < 0$ (see, for example, Henry [8]). Consequently, it is sufficient to show that the hypotheses of the theorem imply $\lambda_1 < 0$ if v is not a constant and this will be the case if we show that $\mathscr{U}(av_X) < 0$ if v is not a constant. We have

$$\mathcal{U}(av_{x}) = \int_{0}^{1} [a[(av_{x})_{x}]^{2} - f'(v)(av_{x})^{2}] dx$$

$$= \int_{0}^{1} af^{2}(v) dx - \int_{0}^{1} f(v)_{x} a av_{x} dx$$

$$= \int_{0}^{1} af^{2}(v) dx + \int_{0}^{1} f(v) a(av_{x})_{x} dx + \int_{0}^{1} f(v) a^{\dagger} av_{x} dx$$

$$= -\int_{0}^{1} (av_{x})_{x} a^{\dagger}(av_{x}) dx$$

To compute the last expression, we integrate by parts to obtain

$$\int_{0}^{1} (av_{x})_{x} a'(av_{x}) dx = -\int_{0}^{1} av_{x} a''av_{x} dx - \int_{0}^{1} av_{x} a'(av_{x})_{x} dx$$

and, thus,

$$\int_0^1 (av_x)_x a'(av_x) dx = -\frac{1}{2} \int_0^1 a''(av_x)^2 dx.$$

Therefore,

(5)
$$\mathscr{U}(av_x) = \frac{1}{2} \int_0^1 a''(av_x)^2 dx \le 0$$

since we have assumed a" < 0.

If $\mathscr{U}(av_X) < 0$, then $\lambda_1 < 0$ and the solution v is unstable. Thus, suppose $\lambda_1 \geq 0$. From (5), this implies $\mathscr{U}(av_X) = 0$ which implies $\lambda_1 = 0$. Since the first eigenvalue of L is simple, there is a ϕ satisfying $L\phi = 0$, $\phi_X = 0$ at x = 0, x = 1, $|\phi|_{L^2(0,1)} = 1$. Also, $\mathscr{U}(av_X) = 0$ implies there is a constant c such that $av_X = c\phi$ on [0,1]. If $c \neq 0$, then $v_X = 0$ at x = 0, x = 1 and a > 0 imply $\phi = 0$ at x = 0, x = 1. But this would imply $\phi = 0$ on [0,1] which is a contradiction. Thus, c = 0 and $av_X = 0$, $av_X = 0$ and $av_X = 0$.

<u>Proof of Theorem 2.</u> Suppose v is an equilibrium solution, $v_{\chi} = 0$ at $x = \alpha$, $x = \beta$, $\alpha, \beta \in (0,1)$. Then v is an equilibrium solution of the Neumann problem on the interval $[\alpha, \beta]$. As in the proof of Theorem 1, one shows that the first eigenvalue λ_1 of the linear variational operator is negative if v is not a constant. By the characterization of λ_1 as a minimum of the functional $\mathcal{U}(u)$, it follows that λ_1 for the Dirichlet problem on [0,1] is negative. This proves the theorem.

REFERENCES

- [1] R.G. Casten and C.J. Holland, Instability results for a reaction diffusion equation with Neumann boundary conditions. J. Differential Equations 27 (1978), 266-273.
- [2] N. Chafee, Asymptotic behavior of a one-dimensional heat equation with homogeneous Neumann boundary conditions. J. Differential Equations 18 (1975), 111-134.
- [3] N. Chafee, The electric ballast resistor: homogeneous and nonhomogeneous equilibria. In Nonlinear Differential equations: Invariance, Stability and Bifurcation. P. de Mottoni and L. Salvadori eds. Academic Press (1981), 161-173.
- [4] N. Chafee and E. F. Infante, A bifurcation problem for a nonlinear partial differential equation of parabolic type. J. Math. Anal. & Appl. 4(1974), 17-37.
- [5] W. Fleming, A selection-migration model in population genetics. J. Math. Biol. 2(1975), 219-233.
- [6] J.K. Hale and P. Massatt, Asymptotic behavior of gradient-like systems. Univ. of Fla. Symp. Dyn. Systems, II, Academic Press, 1982.
- [7] J.K. Hale and J. Vegas, A nonlinear parabolic equation with varying domain. Arch. Rat. Mech. Ana. To appear.
- [8] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Lecture Notes in Math., Vol. 840, Springer-Verlag, 1981.
- [9] B.L. Keyfitz and H.J. Kuiper, Bifurcations resulting from changes in domain in a reaction diffusion equation. J. Differential Eqns. To appear
- [10] K. Maginu, Stability of stationary solutions of one-dimensional semilinear parabolic partial differential equation, J. Math. Anal. Appl. 3(1978), 224-243.
- [11] H. Matano, Convergence of solutions of one-dimensional semilinear parabolic equations. J. Math. Kyoto University 18(1978), 221-227.
- [12] H. Matano, Asymptotic behavior and stability of solutions of semilinear diffusion equations. Res. Inst. Math. Sci., Kyoto 15(1979), 401-454.
- [13] H. Matano, Nonincrease of lap number of a solution for a one dimensional semilinear parabolic equation. Pub. Fac. Sci. Univ. Tokyo. To appear.
- [14] J. Vegas, Bifurcations caused by perturbing the domain in an elliptic equation. J. Differential Eqns. To appear.
- [15] T.I. Zelenyak, Stabilization of solutions of boundary value problems for a second order parabolic equation with one space variable. Differential Equations $\underline{4}(1968)$, 17-22.

FILMED